

## GI Enrollment Form

Phone: (877)229-1724 | Fax: (877)229-1725

## PATIENT INFORMATION PRESCRIBER INFORMATION (Complete the following or send patient demographic sheet.) Please fax copy of patient's insurance card including both sides Prescriber Name: \_\_\_\_\_ State Lic. #: NPI: DOB:\_\_\_\_\_\_ Gender: M / F Height: \_\_\_\_\_ Wt:\_\_\_\_\_ Facility Name: \_\_\_\_\_ City, State, Zip: City, State, Zip: Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Home Phone: Contact: Alternate Phone: Phone: \_\_\_\_\_Email: \_\_\_\_ Last 4 digits SS#:\_\_\_\_\_ Email: \_\_\_\_\_ CLINICAL INFORMATION – STATEMENT OF MEDICAL NECESSITY ☐ K50.90 Crohn's disease NOS ■ K51.90 Ulcerative Colitis ■ Other: Please indicate current or previous treatments and treatment duration below: ■ NSAIDS Duration: \_\_\_\_ Duration: Corticosteroids Duration: Duration: Methotrexate Azathioprine Duration: Sulfasalazine Duration: □ 5 – ASA Other: Duration: Duration: □ 6 – MP Other medications patient is currently taking:\_\_\_\_\_ Date of last dose: Expected date of first/next dose: PRESCRIPTION INFORMATION Medication **Directions** Quantity Refills ☐ Starter Kit 1 starter kit 0 ☐ Initial dose: Inject 400 mg SC at week 0, week 2, and week 4 ☐ Prefilled Syringe Cimzia® ☐ Maintenance dose: Inject 400 mg SC every 4 weeks 28 day supply ☐ Lyophilized Powder Initial dose ☐ Initial Dose: Infuse 300mg IV over 30 minutes at week 0, week 2 and week 6 Entyvio® ☐ 300 mg vial ☐ Maintenance dose: Infuse 300mg IV over 30 minutes every 8 weeks QS ☐ Starter Kit 1 starter kit n ☐ 40 mg Pen ☐ Initial Dose: Inject 160 mg SC on day 1, 80 mg on day 15, then 40 mg ☐ 40 mg Prefilled Syringe thereafter beginning on day 29 Humira® ☐ Inject 40 mg SC every other week ☐ Starter Kit CF 28 day supply ☐ 40 mg CF Pen ☐ Inject 40 mg SC once a week ☐ 40 mg CF Prefilled Syringe ☐ Initial Dose: Infuse \_\_\_\_mg/kg (\_\_\_\_mg) IV at weeks 0, 2, and 6 weeks. Remicade® ☐ 100 mg vial ☐ Maintenance Dose: Infuse \_\_\_\_\_mg/kg (\_\_\_\_mg) IV every \_\_\_\_ weeks. 3 units Ω ☐ 100 mg SmartJect ☐ Initial Dose: Inject 200mg SC at week 0, then 100mg at week 2, then maintenance dose Simponi® ☐ Maintenance Dose: Inject 100mg SC once every 4 weeks ☐ 100 mg Prefilled Syringe 28 day supply **Initial Dose:** Infuse IV over at least 1 hour. $\square \leq 121$ lbs 260 mg (2 vials) ☐ 130 mg vial OS 0 $\Box$ > 121 lbs to 187 lbs 390 mg (3 vials) Stelara® $\square$ > 187 lbs 520 mg (4 vials) ☐ Maintenance Dose: Inject 90 mg SC 8 weeks after initial QS ☐ 90mg Prefilled Syringe IV infusion, then every 8 weeks thereafter Xifaxan® ☐ 550mg Tablet ☐ Take 1 tablet by mouth 3 times a day Other: Ship Medications To: Physician's Clinic Patient's Home ☐ Yes Injection Training Needed: ☐ No \_ Brand Name Required? I authorize this pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorization forms, financial treatment and the receipt and submission of patient lab values and other patient data. In the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize this pharmacy to forward this information and any related materials related to coverage of this product to another pharmacy of the patient's choice or in the